Global Health & Medicine; DOI: 10.35772/ghm.2022.01069

Supplemental Table S1. Current status of long-term sequelae of COVID-19

First author (Ref.)	Study location	Date of admission	Data collection	No.	Timing of follow-up	Prevalence of sequelae	Frequency of specific symptoms
Bai et al. (7)	Milan, Italy	March 1 and November 1, 2020	On-site questionnaire	377	1-3 months after discharge	69%	Fatigue (39.5%), exertional dyspnoea (28.9%)
Huang et al. (11)	Wuhan, China	January 7 and May 29, 2020	On-site questionnaire	1,276	6 months after discharge	76%	Fatigue or muscle weakness (63%), sleep difficulties (26%), hair loss (22%)
Huang et al. (12)	Wuhan, China	Jan 7 and May 29, 2020	On-site questionnaire	1,272	12 months after discharge	49%	Fatigue or muscle weakness (20%), sleep difficulties (17%)
Chen et al. (13)	Wuhan, China	Before April 1, 2020	Telephone interview	715	6 months after discharge	29.9%	Respiratory symptoms (22.7%), fatigue (19.2%),
Shivani et al. (14)	Pakistan	July 2020 to December 2021	Telephone interview	1,498	12 months after discharge		Fatigue (6.78%), dyspnea (4.56%)

No.: Number of patients with COVID-19.